

University of Groningen

Carathéodory and Helly-numbers of convex-product-structures

Sierksma, Gerard

Published in:
Pacific Journal of Mathematics

DOI:
[10.2140/pjm.1975.61.275](https://doi.org/10.2140/pjm.1975.61.275)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1975

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sierksma, G. (1975). Carathéodory and Helly-numbers of convex-product-structures. *Pacific Journal of Mathematics*, 61(1), 275-282. <https://doi.org/10.2140/pjm.1975.61.275>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Pacific Journal of Mathematics

**CARATHÉODORY AND HELLY-NUMBERS OF
CONVEX-PRODUCT-STRUCTURES**

GERARD SIERKSMA

CARATHÉODORY AND HELLY-NUMBERS OF CONVEX-PRODUCT-STRUCTURES

GERARD SIERKSMA

Let c_1 and c_2 be the Carathéodory-numbers of the convexity-structures \mathcal{C}_1 for X_1 , respectively \mathcal{C}_2 for X_2 . It is shown that the Carathéodory-number c of the convex-product-structure $\mathcal{C}_1 \oplus \mathcal{C}_2$ for $X_1 \times X_2$ satisfies the inequality $c_1 + c_2 - 2 \leq c \leq c_1 + c_2$; $c_1, c_2 \geq 2$.

The upper bound for c can be improved by one, resp. two, if a certain number, namely the so-called exchange-number, of one resp. each of the structures \mathcal{C}_1 and \mathcal{C}_2 is less than or equal to the Carathéodory-number of that structure.

A new definition of the Helly-number is given and Levi's theorem is proved with this new definition. Finally it is shown that the Helly-number of a convex-product-structure is the greater of the Helly-numbers of \mathcal{C}_1 and \mathcal{C}_2 .

1. Preliminary remarks and definitions. Existing notations and definitions have been taken from [3], [4] and, in particular, from [8]. Let \mathcal{C} be a collection of subsets of a set X ; by $\bigcap \mathcal{C}$ and $\bigcup \mathcal{C}$ we denote the intersection and the union respectively, of the elements of \mathcal{C} . \mathcal{C} is called a *convexity-structure* for X iff $\emptyset \in \mathcal{C}$, $X \in \mathcal{C}$ and $\bigcap \mathcal{F} \in \mathcal{C}$ for each subcollection $\mathcal{F} \subset \mathcal{C}$; the pair (X, \mathcal{C}) is called a *convexity-space*. The \mathcal{C} -*hull* of a set $S \subset X$, denoted by $\mathcal{C}(S)$, is defined by $\mathcal{C}(S) = \bigcap \{C \mid C \in \mathcal{C} \wedge S \subset C\}$. We shall write $\mathcal{C}(a_1, \dots, a_n)$ instead of $\mathcal{C}(\{a_1, \dots, a_n\})$, and $\mathcal{C}(p \cup (A \setminus a))$ instead of $\mathcal{C}(\{p\} \cup (A \setminus \{a\}))$.

Let X_i be a nonempty set and let \mathcal{C}_i be a convexity-structure for X_i ; $i = 1, 2$. Then $\mathcal{C}_1 \oplus \mathcal{C}_2 = \{A \times B \mid A \in \mathcal{C}_1 \wedge B \in \mathcal{C}_2\}$ is a convexity-structure for the Cartesian-product $X_1 \times X_2$. The pair $(X_1 \times X_2, \mathcal{C}_1 \oplus \mathcal{C}_2)$ is called the *convex-product-space*, also called the *Eckhoff-space*. Note that the $\mathcal{C}_1 \oplus \mathcal{C}_2$ -hull of $E \subset X_1 \times X_2$ is given by $(\mathcal{C}_1 \oplus \mathcal{C}_2)(E) = \mathcal{C}_1(\pi_1 E) \times \mathcal{C}_2(\pi_2 E)$, where π_i is the projection of $X_1 \times X_2$ on X_i ; $i = 1, 2$. Also note that if $e_1, e_2, e_3 \in X_1 \times X_2$ with $e_1 \neq e_2$ and $\pi_i(e_i) = \pi_i(e_3)$ for $i = 1, 2$, then $e_3 \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(e_1, e_2)$.

2. The Carathéodory-number and the exchange-number. A convexity-structure \mathcal{C} for X is said to possess the *Carathéodory-number* c if c is the smallest nonnegative integer such that $\mathcal{C}(S) = \bigcup \{\mathcal{C}(T) \mid T \subset S \wedge |T| \leq c\}$, for all $S \subset X$. The following lemma is an immediate consequence of this definition.

LEMMA 2.1. Let \mathcal{C} be a convexity-structure for X with Carathéodory-number c and let $f \in N$ ($N = 1, 2, 3, \dots$). Then the following holds:

- (i) $(\exists A)[A \subset X \wedge |A| = c \wedge \mathcal{C}(A) \not\subset \bigcup \{\mathcal{C}(A \setminus a) \mid a \in A\}]$;
- (ii) $(\exists A)[A \subset X \wedge |A| = f \wedge \mathcal{C}(A) \not\subset \bigcup \{\mathcal{C}(A \setminus a) \mid a \in A\}] \Rightarrow c \geq f$.

DEFINITION 2.1. The *exchange-number* of a convexity-structure \mathcal{C} for X is the smallest positive integer e , such that

$$(\forall A)(\forall p)[A \subset X \wedge p \in X \wedge e \leq |A| < \infty \\ \implies \mathcal{C}(A) \subset \bigcup \{\mathcal{C}(p \cup (A \setminus a)) \mid a \in A\}].$$

Of course, if the exchange-number e of the convexity-structure \mathcal{C} for X exists then $e \geq 1$; if \mathcal{C} is a T_1 -convexity-structure (see [4]) then $e \geq 2$; if $A \subset X$, $|A| \geq e$ and $p \in \mathcal{C}(A)$ then $\mathcal{C}(A) = \bigcup \{\mathcal{C}(p \cup (A \setminus a)) \mid a \in A\}$, see [5], axiom C3; if the Carathéodory-number c of \mathcal{C} exists too, then $e \leq c + 1$, which follows directly from Lemma 2.1(ii).

EXAMPLE 2.1. Take $X = \mathbf{R}^n$ ($n \in N$) and $\mathcal{C} = \text{conv}$, the usual convexity-structure for the Euclidean-space \mathbf{R}^n . The classical theorem of Carathéodory implies that $c = n + 1$; see [2]. In [6] J. R. Reay proved that $e = n + 1$.

EXAMPLE 2.2. Let $\mathcal{C} = \{X\} \cup \{A \mid A \subset X \wedge |A| \leq f\}$, $f \in N$. Then $c = f + 1$ and $e = 2$.

EXAMPLE 2.3. Take $M \subset X$, $|M| = m$ ($m \in N$), and define $\mathcal{C} = \{X\} \cup \{A \mid A \subset X \wedge M \not\subset A\}$. Because $\mathcal{C}(A) = X$ if $M \subset A$, and $\mathcal{C}(A) = A$ if $M \not\subset A$, it follows that $c = m$ and $e = m + 1$.

EXAMPLE 2.4. The convexity-structure $\mathcal{C} = \{X\} \cup \{A \mid A \subset X \wedge |A| < \infty\}$, with $|X| = \infty$, has no Carathéodory-number, but the exchange-number is 2.

EXAMPLE 2.5. The convex-product-structure $\text{conv} \oplus \text{conv}$ for $\mathbf{R} \times \mathbf{R}$ has Carathéodory-number 2 and exchange-number 3.

THEOREM 2.1. Let \mathcal{C}_i be a convexity-structure for X_i , $X_i \neq \emptyset$; let c_i and e_i be the Carathéodory-number respectively the exchange-number of \mathcal{C}_i ; $i = 1, 2$. The Carathéodory-number c of $\mathcal{C}_1 \oplus \mathcal{C}_2$ exists and the following assertions hold:

- I. If $\min(c_1, c_2) = 1$ then
 - a. $c_1 + c_2 - 1 \leq c \leq c_1 + c_2$

- b. $(\exists i)[i \in \{1, 2\} \wedge e_i \leq c_i] \Rightarrow c = c_1 + c_2 - 1$.
- II. If $\min(c_1, c_2) \geq 2$ then
- a. $c_1 + c_2 - 2 \leq c \leq c_1 + c_2$
- b. $(\exists i)[i \in \{1, 2\} \wedge e_i \leq c_i] \Rightarrow c_1 + c_2 - 2 \leq c \leq c_1 + c_2 - 1$
- c. $(\forall i)[i \in \{1, 2\} \wedge e_i \leq c_i] \Rightarrow c = c_1 + c_2 - 2$.

Proof. First we show that the Carathéodory-number c of $\mathcal{C}_1 \oplus \mathcal{C}_2$ exists and that $c \leq c_1 + c_2$. Let $\emptyset \neq E \subset X_1 \times X_2$ and $(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(E) = \mathcal{C}_1(\pi_1 E) \times \mathcal{C}_2(\pi_2 E)$. Hence there exists a set $S_i \subset \pi_i E$, $|S_i| \leq c_i$ such that $a_i \in \mathcal{C}_i(S_i)$; $i = 1, 2$. $S_i \subset \pi_i E$ implies, there exists a set $F_i \subset E$ such that $\pi_i F_i = S_i$ and $|F_i| = |S_i|$; $i = 1, 2$. So $(a_1, a_2) \in \mathcal{C}_1(S_1) \times \mathcal{C}_2(S_2) = \mathcal{C}_1(\pi_1 F_1) \times \mathcal{C}_2(\pi_2 F_2) \subset \mathcal{C}_1(\pi_1(F_1 \cup F_2)) \times \mathcal{C}_2(\pi_2(F_1 \cup F_2)) = (\mathcal{C}_1 \oplus \mathcal{C}_2)(F_1 \cup F_2)$. Obviously $|F_1 \cup F_2| \leq c_1 + c_2$. Because $(\mathcal{C}_1 \oplus \mathcal{C}_2)(F) \subset (\mathcal{C}_1 \oplus \mathcal{C}_2)(E)$ for each $F \subset E$, we have $(\mathcal{C}_1 \oplus \mathcal{C}_2)(E) = \bigcup \{(\mathcal{C}_1 \oplus \mathcal{C}_2)(F) \mid F \subset E \wedge |F| \leq c_1 + c_2\}$; hence c exists and $c \leq c_1 + c_2$.

To determine the lower bound for c we choose, according to Lemma 2.1.(i), a set $A_i \subset X_i$ such that $|A_i| = c_i$ and $\mathcal{C}_i(A_i) \not\subset \bigcup \{\mathcal{C}_i(A_i \setminus a) \mid a \in A_i\}$; $i = 1, 2$. Take $b_1 \in A_1$ and $b_2 \in A_2$, and consider the set $G = (A_1 \times \{b_2\}) \cup (\{b_1\} \times A_2) \subset X_1 \times X_2$. Obviously $|G| = c_1 + c_2 - 1$. There are two cases (take $c_1 \leq c_2$):

1. Let $c_1 = 1$. Then $A_1 = \{b_1\}$ and $G = \{b_1\} \times A_2$. So we have $(\mathcal{C}_1 \oplus \mathcal{C}_2)(G) = \mathcal{C}_1(b_1) \times \mathcal{C}_2(A_2) \not\subset \mathcal{C}_1(b_1) \times \bigcup \{\mathcal{C}_2(A_2 \setminus b) \mid b \in A_2\} = \bigcup \{(\mathcal{C}_1 \oplus \mathcal{C}_2)(G \setminus e) \mid e \in G\}$, and it follows from Lemma 2.1(ii) that $c \geq c_1 + c_2 - 1$.

2. Let $c_1 \geq 2$. Then also $c_2 \geq 2$. So there exists an element $d_i \in A_i$ with $d_i \neq b_i$; $i = 1, 2$. Note that $\pi_1((b_1, b_2)) = \pi_1((b_1, d_2))$ and $\pi_2((b_1, b_2)) = \pi_2((d_1, b_2))$. The last remark of §1 gives us that $(b_1, b_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)((d_1, b_2), (b_1, d_2)) \subset (\mathcal{C}_1 \oplus \mathcal{C}_2)(G \setminus (b_1, b_2))$. Define $F = G \setminus \{(b_1, b_2)\}$. Clearly $|F| = c_1 + c_2 - 2$ and $\pi_i F = A_i$; $i = 1, 2$. Moreover $(\mathcal{C}_1 \oplus \mathcal{C}_2)(F) \not\subset \bigcup \{(\mathcal{C}_1 \oplus \mathcal{C}_2)(F \setminus e) \mid e \in F\}$. From Lemma 2.1(ii) it follows that $c \geq c_1 + c_2 - 2$.

We now prove I.b and II.b. Let us assume that $e_i \leq c_i$. Take $c_1, c_2 \geq 1$, $\emptyset \neq E \subset X_1 \times X_2$ and $(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(E)$. We show that there exists a set F such that

$$(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(F), \quad F \subset E \quad \text{and} \quad |F| \leq c_1 + c_2 - 1.$$

The first part of the proof of this theorem implies that there exists a set F_i such that $a_i \in \mathcal{C}_i(\pi_i F_i)$, $F_i \subset E$ and $|\pi_i F_i| = |F_i| \leq c_i$; $i = 1, 2$. We may assume that $|F_i| = c_i \quad \forall i \in \{1, 2\}$, because if $|F_i| \leq c_i - 1$ for some $i \in \{1, 2\}$ then $|F_1 \cup F_2| \leq c_1 + c_2 - 1$, and we are done. If $F_1 \cap F_2 \neq \emptyset$, then define $F = F_1 \cup F_2$. So $|F| \leq c_1 + c_2 - 1$ and $(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(F)$. If $F_1 \cap F_2 = \emptyset$, we distinguish two cases:

1. $(\exists i)[i \in \{1, 2\} \wedge \pi_i F_1 \cap \pi_i F_2 \neq \emptyset]$. Take $i = 1$. Hence $\pi_1 F_1 \cap \pi_1 F_2 \neq \emptyset$. Now there exist elements $e_1 \in F_1$ and $e_2 \in F_2$ such that $\pi_1(e_1) = \pi_1(e_2)$. Note that $e_1 \neq e_2$. Define $F = (F_1 \setminus \{e_1\}) \cup F_2$. Clearly $|F| \leq c_1 + c_2 - 1$ and $\pi_i F_i \subset \pi_i F$; $i = 1, 2$. So $(a_1, a_2) \in \mathcal{C}_1(\pi_1 F_1) \times \mathcal{C}_2(\pi_2 F_2) \subset \mathcal{C}_1(\pi_1 F) \times \mathcal{C}_2(\pi_2 F) = (\mathcal{C}_1 \oplus \mathcal{C}_2)(F)$.

2. $(\forall i)[i \in \{1, 2\} \Rightarrow \pi_i F_1 \cap \pi_i F_2 = \emptyset]$. Take $e \in F_2$. Then $\pi_1(e) \notin \pi_1 F_1$. Because $c_1 \leq c_1$, there exists an element $e_1 \in F_1$ such that $a_1 \in \mathcal{C}_1(\pi_1(e) \cup \pi_1 F_1 \setminus \pi_1(e_1)) \subset \mathcal{C}_1(\pi_1(e \cup F_1 \setminus \{e_1\}))$. Define $F = (F_1 \setminus \{e_1\}) \cup F_2$. Obviously $|F| \leq c_1 + c_2 - 1$, $a_1 \in \mathcal{C}_1(\pi_1 F)$ and $a_2 \in \mathcal{C}_2(\pi_2 F_2) \subset \mathcal{C}_2(\pi_2 F)$. Hence $(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(F)$.

Finally we prove II.c. Take again $\emptyset \neq E \subset X_1 \times X_2$ and $(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(E)$. We shall show that there exists a set F such that

$$(a_1, a_2) \in (\mathcal{C}_1 \oplus \mathcal{C}_2)(F), \quad F \subset E, \quad \text{and} \quad |F| \leq c_1 + c_2 - 2.$$

In the proof of II.b we found a set $G_i \subset E$ such that $|\pi_i G_i| = |G_i| \leq c_i$, $a_i \in \mathcal{C}(\pi_i G_i)$ and $|G_1 \cup G_2| \leq c_1 + c_2 - 1$; $i = 1, 2$. As in the proof of II.b we may assume that $|G_i| = c_i$; $i = 1, 2$. If $|G_1 \cup G_2| > 1$ then define $F = G_1 \cup G_2$, hence $a_i \in \mathcal{C}_i(\pi_i F)$ and $|F| \leq c_1 + c_2 - 2$, so we are done. The case that $|G_1 \cap G_2| = 1$ still remains. Assume $G_1 \cap G_2 = \{e\}$, and $|G_i| \geq 2$ for $i = 1, 2$. Throughout the remainder of the proof we take $i, j \in \{1, 2\}$ with $i + j = 3$. Let $e_i \in G_j$, $e_i \neq e$. There are two cases:

1. If $\pi_i(e_i) \notin \pi_i G_i$ then, because $c_i \leq c_i$, there exists an element $u_i \in G_i$ such that $u_i \neq e_i$ and $a_i \in \mathcal{C}_i(\pi_i(e_i) \cup (\pi_i G_i \setminus \pi_i(u_i))) \subset \mathcal{C}_i(\pi_i(e_i \cup (G_i \setminus u_i)))$.

2. If $\pi_i(e_i) \in \pi_i G_i$ then there exists an element $v_i \in G_i$ such that $v_i \neq e_i$ and $\pi_i(e_i) = \pi_i(v_i)$; hence $\pi_i G_i = \pi_i(e_i \cup (G_i \setminus v_i))$, so $a_i \in \mathcal{C}_i(\pi_i G_i) = \mathcal{C}_i(\pi_i(e_i \cup (G_i \setminus v_i)))$.

We may conclude that in both cases there exists an element $d_i \in G_i$ such that $d_i \neq e_i$ and $a_i \in \mathcal{C}_i(\pi_i(e_i \cup (G_i \setminus d_i)))$.

If $d_1 = e = d_2$ then define $F = G_1 \cup G_2 \setminus \{e\}$. Hence $|F| \leq c_1 + c_2 - 2$ and, because $e_i \in G_j$, $e_i \neq e$ we have $(a_1, a_2) \in \mathcal{C}_1(\pi_1(e_1 \cup (G_1 \setminus e))) \times \mathcal{C}_2(\pi_2(e_2 \cup (G_2 \setminus e))) \subset (\mathcal{C}_1 \oplus \mathcal{C}_2)(G_1 \cup G_2 \setminus e) = (\mathcal{C}_1 \oplus \mathcal{C}_2)(F)$. If $d_i \neq e$ for some $i \in \{1, 2\}$ then, taking e.g. $i = 1$, we define $F = G_1 \cup G_2 \setminus \{d_1\}$. Clearly $|F| \leq c_1 + c_2 - 2$ and because $e_1 \in G_2$ we have $(a_1, a_2) \in \mathcal{C}_1(\pi_1(e_1 \cup (G_1 \setminus d_1))) \times \mathcal{C}_2(\pi_2 G_2) \subset (\mathcal{C}_1 \oplus \mathcal{C}_2)(G_1 \cup G_2 \setminus d_1) = (\mathcal{C}_1 \oplus \mathcal{C}_2)(F)$.

It follows that $c \leq c_1 + c_2 - 2$ and because $c \geq c_1 + c_2 - 2$, as we have seen already, we may conclude that $c = c_1 + c_2 - 2$. This completes the proof of Theorem 2.1.

EXAMPLE 2.6. Take $X_1 = \mathbf{R}^m$ and $X_2 = \mathbf{R}^n$ ($m, n \in \mathbf{N}$), and $\mathcal{C}_1 = \text{conv} = \mathcal{C}_2$ (see Example 2.1). Because $c_1 = e_1 = m + 1$ and $c_2 = e_2 = n + 1$ it follows from Theorem 2.1. (II.c) that the Carathéodory-number of the convex-product-structure $\text{conv} \oplus \text{conv}$ for \mathbf{R}^{m+n} is $c = c_1 + c_2 - 2 = m + n$.

EXAMPLE 2.7. Take $\mathcal{C}_i = \{X_i\} \cup \{A \mid A \subset X_i \wedge |A| \leq f_i\}$, $f_i \geq 1$, then because $e_i = 2 \leq f_i + 1 = c_i$, $\forall i \in \{1, 2\}$, it follows from Theorem 2.1(II.c) that $c = c_1 + c_2 - 2 = f_1 + f_2$.

EXAMPLE 2.8. Take $M_i \subset X_i$, $|M_i| = m_i$ and define $\mathcal{C}_i = \{X_i\} \cup \{A \mid A \subset X_i \wedge M_i \not\subset A\}$; $i = 1, 2$. Because $e_i = m_i + 1 > m_i = c_i$ it follows from Theorem 2.1(I.II.a) that $c \leq c_1 + c_2 = m_1 + m_2$. We shall show now that even $c = m_1 + m_2$. Consider a set $E \subset X_1 \times X_2$ such that $E = E_1 \cup E_2$, with $|E_i| = m_i$, $|\pi_i E_i| = 1$, $\pi_i E_j = M_i$ and $\pi_i E_1 \cap \pi_i E_2 = \emptyset$; $i, j \in \{1, 2\}$ and $i + j = 3$. It is easy to see that $(\mathcal{C}_1 \oplus \mathcal{C}_2)(E) = X_1 \times X_2$. However, in general, $\bigcup \{(\mathcal{C}_1 \oplus \mathcal{C}_2)(E \setminus e) \mid e \in E\} = (\pi_1 E \times X_2) \cup (X_1 \times \pi_2 E) \neq X_1 \times X_2$. Hence $c = m_1 + m_2$.

EXAMPLE 2.9. Take $M \subset X_1$, $|M| = m$ and $\mathcal{C}_1 = \{X_1\} \cup \{A \mid A \subset X_1 \wedge M \not\subset A\}$. Take $X_2 = \mathbf{R}^n$ and $\mathcal{C}_2 = \text{conv}$. We know that $e_1 = m + 1 > m = c_1$, and $e_2 = n + 1 = c_2$. From Theorem 2.1(II.b) it follows that $c \leq c_1 + c_2 - 1 = m + n$. As in the previous example we can show that $c = m + n$. In order to prove this we have to look for a set $E \subset X_1 \times X_2$, $|E| = m + n$, such that the convex hull of E is not the union of the convex hulls of proper subsets of E . Take $E = E_1 \cup E_2$ with $|E_1| = m$, $|E_2| = n$, $|\pi_i E_j| = 1$, $\pi_i E_1 \cap \pi_i E_2 = \emptyset$, $\pi_1 E_1 = M$ and $\mathcal{C}_2(\pi_2 E) \neq \bigcup \{\mathcal{C}_2(\pi_2 E \setminus a) \mid a \in \pi_2 E\}$; $i, j \in \{1, 2\}$, $i + j = 3$. Note that $|\pi_2 E| = n + 1$. Now we have $(\mathcal{C}_1 \oplus \mathcal{C}_2)(E) = X_1 \times \mathcal{C}_2(\pi_2 E)$. But, in general, $\bigcup \{(\mathcal{C}_1 \oplus \mathcal{C}_2)(E \setminus e) \mid e \in E\} \neq X_1 \times \mathcal{C}_2(\pi_2 E)$. Hence $c = m + n$.

The main result of this section, Theorem 2.1, is a generalization of J. R. Reay's Theorem 1 (first part) together with Example (1) on pg. 229 of [7]. In fact, Reay proves that the Carathéodory-number c of the convex-product-structure, whose component spaces are Euclidian-spaces with dimensions m_1 and n_2 and with the usual convexity-structure, satisfies the condition $c = m_1 + n_2$. (See Ex. 2.6.)

3. The Helly-number. Let $\mathcal{F} \subset 2^X$, $|F| \geq f$, $f \in \mathbf{N}$, for some set X . Define $\bigcap_{(f)} F = \{\bigcap A \mid A \in \mathcal{F}^f\}$. I.e. $\emptyset \notin \bigcap_{(f)} \mathcal{F}$ implies that each intersection of f elements of \mathcal{F} is not empty.

DEFINITION 3.1. A convexity-structure \mathcal{C} for X has the *Helly-number* h if h is the smallest integer such that $[\mathcal{F} \subset \mathcal{C} \wedge$

$$|\mathcal{F}| < \infty \wedge \emptyset \notin \bigcap_{(t)} F] \Rightarrow \bigcap \mathcal{F} \neq \emptyset.$$

Note that if $X = \emptyset$ then \mathfrak{h} does not exist, and if \mathfrak{h} exists then $\mathfrak{h} \geq 1$. The following characterization of the Helly-number gives rise to another definition of it; see also [1] and [4].

LEMMA 3.1. *Let \mathcal{C} be a convexity-structure for X with Helly-number \mathfrak{h} . Then the following assertions are equivalent:*

- (i) $\mathfrak{h} \leq \mathfrak{k}$;
- (ii) $[\mathcal{F} \subset \mathcal{C} \wedge |\mathcal{F}| = \mathfrak{k} + 1 \wedge \emptyset \notin \bigcap_{(t)} \mathcal{F}] \Rightarrow \bigcap \mathcal{F} \neq \emptyset$;
- (iii) $[\mathcal{A} \subset X \wedge |A| = \mathfrak{k} + 1] \Rightarrow \bigcap \{\mathcal{C}(A \setminus a) \mid a \in A\} \neq \emptyset$;
- (iv) $[A \subset X \wedge \mathfrak{k} + 1 \leq |A| < \infty] \Rightarrow \bigcap \{\mathcal{C}(A \setminus a) \mid a \in A\} \neq \emptyset$.

Proof. We shall go through the following implication-cycle:

(i) \Rightarrow (iv) \Rightarrow (iii) \Rightarrow (ii) \Rightarrow (i).

(i) \Rightarrow (iv): Take $A \subset X$, $|A| = \mathfrak{k} + n$ ($n \in \mathbb{N}$) and define $\mathcal{F} = \{\mathcal{C}(A \setminus a) \mid a \in A\}$. Note that $|\mathcal{F}| \leq \mathfrak{k} + n$. If there exist elements $a, b \in A$, such that $a \neq b$ and $\mathcal{C}(A \setminus a) = \mathcal{C}(A \setminus b)$ then of course $\bigcap \{\mathcal{C}(A \setminus a) \mid a \in A\} \neq \emptyset$, and we are done. So we may assume that $|\mathcal{F}| = \mathfrak{k} + n$. If $\mathcal{F}' \subset \mathcal{F}$, with $|\mathcal{F}'| = \mathfrak{k}$, then $\mathcal{F}' \neq \mathcal{F}$ and there exists an element $a_1 \in A$ such that $\mathcal{C}(A \setminus a_1) \notin \mathcal{F}'$. From $a_1 \in A \setminus \{a\} \subset \mathcal{C}(A \setminus a)$ for each $a \in A$, $a \neq a_1$, it follows that $a_1 \in \bigcap \mathcal{F}'$, and hence $\emptyset \notin \bigcap_{(t)} \mathcal{F}$. Because $\mathfrak{h} \leq \mathfrak{k}$ we have $[\emptyset \notin \bigcap_{(t)} \mathcal{F} \Rightarrow \emptyset \notin \bigcap_{(t)} \mathcal{F}]$. From Definition 3.1 it follows that $\bigcap \{\mathcal{C}(A \setminus a) \mid a \in A\} = \bigcap \mathcal{F} \neq \emptyset$.

(iv) \Rightarrow (iii) is trivial and (ii) \Rightarrow (i) follows by induction. It remains to be shown that (iii) \Rightarrow (ii): Take $\mathcal{F} \subset \mathcal{C}$, $|F| = \mathfrak{k} + 1$ and $\emptyset \notin \bigcap_{(t)} \mathcal{F}$. Let $\mathcal{F} = \{F_i \mid i = 1, 2, \dots, \mathfrak{k} + 1\}$. Then $\emptyset \notin \bigcap_{(t)} \mathcal{F} \Rightarrow \bigcap \{\mathcal{F} \setminus \{F_i\}\} \neq \emptyset$, $\forall i = 1, 2, \dots, \mathfrak{k} + 1$. For each $i = 1, 2, \dots, \mathfrak{k} + 1$ we choose an element $a_i \in \bigcap \{\mathcal{F} \setminus \{F_i\}\}$. Define $A = \{a_i \mid i = 1, 2, \dots, \mathfrak{k} + 1\}$. Hence $|A| \leq \mathfrak{k} + 1$. If $|A| < \mathfrak{k} + 1$, then there exists an index i such that $a_i \in F_i$. But then $a_i \in \bigcap \mathcal{F}$. Hence $\bigcap \mathcal{F} \neq \emptyset$, and we are done. So we may assume that $|A| = \mathfrak{k} + 1$. From (iii) it follows that $\bigcap \{\mathcal{C}(A \setminus a_i) \mid i = 1, 2, \dots, \mathfrak{k} + 1\} \neq \emptyset$. For each $i = 1, 2, \dots, \mathfrak{k} + 1$ we have $A \setminus \{a_i\} \subset F_i$, so $\mathcal{C}(A \setminus a_i) \subset F_i$. Hence $\bigcap \{\mathcal{C}(A \setminus a_i) \mid i = 1, 2, \dots, \mathfrak{k} + 1\} \subset \bigcap \mathcal{F}$, and we conclude that $\bigcap \mathcal{F} \neq \emptyset$.

DEFINITION 3.1'. A convexity-structure \mathcal{C} for X has the *Helly-number* \mathfrak{h} if \mathfrak{h} is the smallest integer such that $[A \subset X \wedge |A| = \mathfrak{h} + 1] \Rightarrow \bigcap \{\mathcal{C}(A \setminus a) \mid a \in A\} \neq \emptyset$.

With the aid of Lemma 3.1. it is easy to verify that Definitions 3.1 and 3.1' are equivalent. We now prove the classical theorem of Levi with the aid of Definition 3.1'. See [5], Theorem H.

THEOREM 3.1 (Levi). *Let \mathcal{C} be a convexity-structure for X . Then the existence of a Radon-number r implies the existence of a Helly-number h , such that $h \leq r - 1$.*

Proof. Each $A \subset X$ with $|A| = r$ has a \mathcal{C} -Radon-partition; see [2]. So there exists a set $B \subset A$, with $\emptyset \neq B \neq A$ and $\mathcal{C}(B) \cap \mathcal{C}(A \setminus B) \neq \emptyset$. Because $\mathcal{C}(B) \subset \bigcap \{\mathcal{C}(A \setminus a) \mid a \in A \setminus B\}$ and $\mathcal{C}(A \setminus B) \subset \bigcap \{\mathcal{C}(A \setminus a) \mid a \in B\}$, we have $\bigcap \{\mathcal{C}(A \setminus a) \mid a \in A\} \supset \mathcal{C}(B) \cap \mathcal{C}(A \setminus B) \neq \emptyset$. Hence h exists and $h \leq r - 1$.

THEOREM 3.2. *Let \mathcal{C}_i be a convexity-structure for X_i , $X_i \neq \emptyset$, with Helly-number h_i ; $i = 1, 2$. Then the Helly-number h of the convex-product-structure $\mathcal{C}_1 \oplus \mathcal{C}_2$ for $X_1 \times X_2$ exists and $h = \max(h_1, h_2)$.*

Proof. We may assume that $h_1 \geq h_2$. Take $E \subset X_1 \times X_2$ with $|E| = h_1 + 1$. There are two possibilities for the projection $\pi_i E$ of E on X_i ; $i = 1, 2$.

—If $|\pi_i E| \leq h_i$ for some $i \in \{1, 2\}$, then there exist elements $e_1, e_2 \in E$ such that $e_1 \neq e_2$ and such that $\pi_i(e_1) = \pi_i(e_2)$. Clearly $\pi_i(e_1) \in \mathcal{C}_i(\pi_i(E \setminus e_1))$. Hence $\pi_i(e_1) \in \bigcap \{\mathcal{C}_i(\pi_i(E \setminus e)) \mid e \in E\} \neq \emptyset$.

—If $|\pi_i E| = h_i + 1$, for some $i \in \{1, 2\}$, then it follows from Definition 3.1' that $\bigcap \{\mathcal{C}_i(\pi_i E \setminus x) \mid x \in \pi_i E\} \neq \emptyset$. Hence $\bigcap \{\mathcal{C}_i(\pi_i(E \setminus e)) \mid e \in E\} \supset \bigcap \{\mathcal{C}_i(\pi_i E \setminus \pi_i(e)) \mid e \in E\} \neq \emptyset$ and we may conclude that $\bigcap \{\mathcal{C}_i(\pi_i(E \setminus e)) \mid e \in E\} \neq \emptyset$; $i = 1, 2$.

Hence $\bigcap \{(\mathcal{C}_1 \oplus \mathcal{C}_2)(E \setminus e) \mid e \in E\} = \bigcap \{\mathcal{C}_1(\pi_1(E \setminus e)) \times \mathcal{C}_2(\pi_2(E \setminus e)) \mid e \in E\} \neq \emptyset$, and so $h \leq h_1 = \max(h_1, h_2)$.

Next we show that $h \geq h_1$. Assuming $h = h_1 - 1$, $A \subset X_1$ with $|A| = h_1$ and $b \in X_2$ we have, because $|A \times \{b\}| = h_1$, $\bigcap \{(\mathcal{C}_1 \oplus \mathcal{C}_2)((A \times \{b\}) \setminus (a, b)) \mid a \in A\} \neq \emptyset$. Hence $\bigcap \{\mathcal{C}_1(A \setminus a) \mid a \in A\} = \bigcap \{\mathcal{C}_1(\pi_1((A \times \{b\}) \setminus (a, b))) \mid a \in A\} \neq \emptyset$. This contradicts the fact that h_1 is the Helly-number of \mathcal{C}_1 , so that, indeed, $h \geq h_1 = \max(h_1, h_2)$ and the final conclusion is that $h = h_1 = \max(h_1, h_2)$.

It is well known that the Carathéodory-number c , the Helly-number h and the Radon-number r of the usual convexity-structure conv for \mathbb{R}^n satisfy the equalities $c = h = r - 1$ ($= n + 1$), that is, in Levi's theorem equality holds. There are however convexity-structures \mathcal{C} where the equality does not hold. This is even the case when \mathcal{C} is a convex-product-structure:

We know that (see also [8]):

$$\begin{aligned} c_1 + c_2 - 2 &\leq c \leq c_1 + c_2 \\ h &= \max(h_1, h_2) \\ \max(r_1, r_2) &\leq r \leq r_1 + r_2 - 2; \end{aligned}$$

c_i , h_i and r_i are resp. the Carathéodory-, Helly- and Radon-number of \mathcal{C}_i ; $i = 1, 2$.

If $c_i = h_i = r_i - 1$, $i = 1, 2$, then we have:

a. if $c_i \geq 3$ (e.g. when \mathcal{C}_i is T_1) then $c \neq h$.

b. if $c = c_1 + c_2$, then $c > r - 1$.

if the exchange-number, $e_i \leq c_i$ and if $r = r_1 + r_2 - 2$ (e.g. in the case $X_i = \mathbf{R}^{i-1}$, $\mathcal{C}_i = \text{conv}$) then $c \leq r - 1$.

c. if $r > \max(r_1, r_2)$ (e.g. in the same case as in b) then $h < r - 1$.

The results in this paper can be extended to convex-product-structures which are products of finitely many convexity-structures.

In a next paper we shall pay more attention to the properties of the exchange number. For example we shall show that under certain conditions, the exchange-number of a convex-product-structure exists and how it can be derived.

ACKNOWLEDGMENT. The author would like to thank Dr. J. Ch. Boland for his many helpful suggestions in the writing of this paper.

REFERENCES

1. J. R. Calder, *Some elementary properties of interval convexities*, J. London Math. Soc., (2), **3** (1971), 422-428.
2. L. Danzer, B. Grünbaum, and V. Klee, *Helly's theorems and its relatives*, Proc. of Symp. in Pure Math., **7** (1963), "Convexity", Amer. Math. Soc., 101-180.
3. J. Eckhoff, *Der Satz von Radon in konvexen Produktstrukturen I*, Monatshefte für Math., **72** (1968), 303-314.
4. D. C. Kay and E. W. Womble, *Axiomatic convexity theory and relationships between the Carathéodory, Helly, and Radon numbers*, Pacific J. Math., **38** (1971), 471-485.
5. F. W. Levi, *On Helly's theorem and the axioms of convexity*, J. Indian Math. Soc., **15** (1951), 65-76.
6. J. R. Reay, *Generalizations of a theorem of Carathéodory*, Memoirs of the American Math. Soc., Nr. **54** (1965).
7. ———, *Carathéodory theorems in convex product structures*, Pacific J. Math., **35** (1970), 227-230.
8. G. Sierksma and J. Ch. Boland, *The least upper bound for the Radon number of an Eckhoff space*, University of Groningen, Econometric Institute, Report 1974.

Received January 2, 1975.

UNIVERSITY OF GRONINGEN, NETHERLANDS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 61, No. 1

November, 1975

Jiří Adámek, V. Koubek and Věra Trnková, <i>Sums of Boolean spaces represent every group</i>	1
Richard Neal Ball, <i>Full convex l-subgroups and the existence of a^*-closures of lattice ordered groups</i>	7
Joseph Becker, <i>Normal hypersurfaces</i>	17
Gerald A. Beer, <i>Starshaped sets and the Hausdorff metric</i>	21
Dennis Dale Berkey and Alan Cecil Lazer, <i>Linear differential systems with measurable coefficients</i>	29
Harald Boehme, <i>Glättungen von Abbildungen 3-dimensionaler Mannigfaltigkeiten</i>	45
Stephen LaVern Campbell, <i>Linear operators for which T^*T and $T + T^*$ commute</i>	53
H. P. Dikshit and Arun Kumar, <i>Absolute summability of Fourier series with factors</i>	59
Andrew George Earnest and John Sollion Hsia, <i>Spinor norms of local integral rotations. II</i>	71
Erik Maurice Ellentuck, <i>Semigroups, Horn sentences and isolic structures</i>	87
Ingrid Fotino, <i>Generalized convolution ring of arithmetic functions</i>	103
Michael Randy Gabel, <i>Lower bounds on the stable range of polynomial rings</i>	117
Fergus John Gaines, <i>Kato-Taussky-Wielandt commutator relations and characteristic curves</i>	121
Theodore William Gamelin, <i>The polynomial hulls of certain subsets of C^2</i>	129
R. J. Gazik and Darrell Conley Kent, <i>Coarse uniform convergence spaces</i>	143
Paul R. Goodey, <i>A note on starshaped sets</i>	151
Eloise A. Hamann, <i>On power-invariance</i>	153
M. Jayachandran and M. Rajagopalan, <i>Scattered compactification for $N \cup \{P\}$</i>	161
V. Karunakaran, <i>Certain classes of regular univalent functions</i>	173
John Cronan Kieffer, <i>A ratio limit theorem for a strongly subadditive set function in a locally compact amenable group</i>	183
Siu Kwong Lo and Harald G. Niederreiter, <i>Banach-Buck measure, density, and uniform distribution in rings of algebraic integers</i>	191
Harold W. Martin, <i>Contractibility of topological spaces onto metric spaces</i>	209
Harold W. Martin, <i>Local connectedness in developable spaces</i>	219
A. Meir and John W. Moon, <i>Relations between packing and covering numbers of a tree</i>	225
Hiroshi Mori, <i>Notes on stable currents</i>	235
Donald J. Newman and I. J. Schoenberg, <i>Splines and the logarithmic function</i>	241
M. Ann Piech, <i>Locality of the number of particles operator</i>	259
Fred Richman, <i>The constructive theory of KT-modules</i>	263
Gerard Sierksma, <i>Carathéodory and Helly-numbers of convex-product-structures</i>	275
Raymond Earl Smithson, <i>Subcontinuity for multifunctions</i>	283
Gary Roy Spoor, <i>Differentiability conditions and bounds on singular points</i>	289
Rosario Strano, <i>Azumaya algebras over Hensel rings</i>	295